

## THE LOCAL DISTRIBUTION OF MONSOON RAINFALL.

By GILBERT T. WALKER.

[Abstract reprinted from *Memoirs of the Indian Meteorological Dept.*, Vol. XXIII, Part II, Calcutta, 1922.]

Experience of the past 13 years shows that the formula of 1908 for forecasting the whole monsoon rainfall of India affords a useful indication in years when the calculated departure is 1" or over, which, on the average, form about half of the number of years. Investigation of the geographical distribution of rainfall leads to a subdivision of India into large homogeneous areas such as northwest India and the Peninsula, for which formulæ have been worked out, giving the total monsoon rainfall and that of August and September. The correlation coefficients range from 0.57 to 0.73. Similar formulæ have also been developed for Upper Burma, Mysore, and Malabar.

THE CALCULATION OF THE DEGREE OF CONTINENTALITY.<sup>1</sup>

By LADISLAS GORCZYŃSKI.

[Abstracted from *Geografiska Annaler*, 1920, H. 4, pp. 324-331.]

The degree of continentality of a given station may be computed by means of the formula

$$k = 1.7 (A - 12 \sin \phi) / \sin \phi \\ = (1.7 A / \sin \phi) - 20.4$$

in which  $k$  is the thermal degree of continentality in per cent,  $A$  the amplitude of the mean annual temperature fluctuation and  $\phi$  the latitude. The expression  $A = 12 \sin \phi$  is found to be in good accord with observations over the oceans. The constant, 1.7, is derived from the assumption that Werchojansk, in eastern Siberia, is representative of 100 per cent continentality. The amplitude of the annual temperature variation is  $65.9^\circ \text{C.}$ , and the latitude is  $67^\circ 33' \text{N.}$

Upon the basis of this formula, the author presents the geographical distribution of continentality, of which three degrees are distinguished:

- (1) Transitional maritime, in which  $k = 0$  to 33 per cent.
- (2) Continental, in which  $k = 34$  to 66 per cent.
- (3) Extreme continental, in which  $k = 67$  to 100 per cent.

One advantage of discussing continentality from the point of view of this formula, as opposed to the direct use of the amplitude of the annual variation of temperature, is shown by an illustration. Considering only amplitude of temperature variation, the Sahara Desert would fall in the same class as western Europe; whereas, computing continentality from this equation, the Sahara lies in groups (2) and (3) above, while western Europe lies in (1).

The values thus calculated show resemblances both to the "isocontinentals" of Zenker and the "equidistants" of Rohrbach, but possess certain advantages over these earlier attempts at discussing continentality.

The mean values of  $k$  (per cent) for the several continents are:

Europe.....	30 (max. 50, Oural).
Asia.....	58 (max. 100, Werchojansk).
Africa.....	36 (max. 75, Sahara).
North America with	
Greenland.....	44 (max. 70, $\phi = 66^\circ \text{N.}$ ).
South America.....	18 (max. 31, $\phi = 30^\circ \text{S.}$ ).
Australia.....	32 (max. 55, interior).
Arctic Zone.....	40 (North Pole, 48).
Antarctic Zone.....	20 (South Pole, 26).

For the Northern Hemisphere (continents and oceans) the value is 20 per cent; for the Southern Hemisphere, 3 per cent.

A comparison of the relation  $A = 12 \sin \phi$  with observed values over extensive ocean areas shows computed values in close agreement with observation. In the Southern Hemisphere,  $A = 9 \sin \phi$  seems to give better agreement. This, of course, would lead to a somewhat different value of  $k$ , but a series of comparisons demonstrates that the differences would not exceed 3 per cent, which is considered sufficiently small.

The author presents world maps of the distribution of continentality as computed by this formula and of the distribution of amplitude of annual temperature variation.—C. L. M.

## BIBLIOGRAPHY.

## RECENT ADDITIONS TO THE WEATHER BUREAU LIBRARY.

C. FITZHUGH TALMAN, Meteorologist in Charge of Library.

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

Bigelow, Frank H.

La teoría de las dos órbitas para la explicación del origen de la radiación. Buenos Aires. 1921. 19 p. figs. 25½ cm. (Anales de la Sociedad científica Argentina. T. 92.)

Brazier, Charles E.

Recherches expérimentales sur les moulinets anémométriques. [Paris. 1920.] p. 157-300. illus. 33 cm. [Extr.: Annales du Bureau central météorologique de France. Année 1914. I. Mémoires. Paris. 1920.]

Froc, Louis.

L'atmosphère en extrême-orient. . . . pt. 1. Les six mois froids, son état normal, ses perturbations. p. 117-198. illus. plates. pt. 2. Les six mois chauds, son état normal, ses perturbations. p. 169-238. illus. [Paris.] 25 cm. [Extr.: Annales hydrographiques. 2 sér. v. 23-24. 1901-1902.]

Gockel, Albert.

Elektrische Strömungen in der Atmosphäre. . . . Freiburg. 1921. 21 p. 23 cm.

Herrmann, C. F. von.

Economic value of Georgia meteorology. Atlanta. 1922. unp. 27 cm. [Christian index, v. 102, nos. 19-20, May, 1922.]

Jardine, James T., &amp; Forsling, Clarence L.

Range and cattle management during drought. Washington. 1922. 84 p. figs. plates. 23 cm. (U. S. Department of agriculture. Bulletin no. 1031.)

Köhler, Hilding.

Zur Kondensation des Wasserdampfes in der Atmosphäre. Kristiania. 1922. 2<sup>e</sup> Mitt. 53 p. illus. plate. 31 cm. (Geofysiske Kommission. Geofysiske Publikationer. v. 2. no. 6.)

Lucanus, Friedrich von.

Die Rätsel des Vogelzuges. Ihre Lösung auf experimentellem Wege durch Aeronautik, Aviatik und Vogelberingung. Langensalza. 1922. viii, 226 p. illus. plate. 24½ cm.

Mikolaschek, Karl.

Die Niederschlags- und Abflussverhältnisse im Egergebiete. n. p. n. d. 38 p. charts. 24½ cm.

Nedelkovitch, Milan.

Exposé sur l'observatoire central de Belgrade. Belgrade. 1909. 15 p. 33 cm.

Schmucker, Matthäus.

Wetter-Kalender für das Jahr 1922 und zwar für jeden Tag. . . . München. [1921.] 6. Jahrg. 182 p. illus. 15½ cm.